# Introduction to Queen Rearing

Miller Method & Demaree Method

30<sup>th</sup> January 2019 - 7:30pm - Anne Rowberry & Dan Job



#### Introduction

- Overview of Queen Rearing
- Miller Method Dedicated Queen rearing method
  - A Few simple manipulations.
- Demaree Method Swarm prevention using bood relocation
  Can be used to raise queens.
- Queen Mating Options



# Queen Rearing – What's Required?

- Patients & perseverance
- Set up & equipment
- Hive manipulations
- Planning & timing
- Queen mating plan





## Why actively rear Queens?

- Selection Raise queens from stock with desired characteristics
  - ➤ Honey production
  - ➤ Temperament
  - > Hygienic behaviour
  - > Efficient use of winter stores
  - > Few signs of disease
  - ➤ Colony growth / Brood production
  - ➤ Low swarming tendency
- Avoid buying in queens / Replace failing queens
- Be sustainable Replace winter losses / Expand colony numbers






#### How the bees do it!

- Cycle 3 / 5 / 7 ish (Egg / Larvae / Pupa)
  - ➤ Don't forget this can vary slightly dependent on temperature
- Why will bees raise new queens
  - ➤ Emergency Convert eggs intended for workers
  - ➤ Swarming Planned (reproduction)
  - ➤ Supersedure Planned (replacement)






- Prepare the frame
  - ➤ Saw tooth ease of subsequent harvesting of QC's
  - > 2-3 spikes 5-10 QC's
- Foundation or drawn comb
- Insert into parent colony
  - ➤ In centre of brood cluster
  - Check at 3 days intervals





- Prepare the frame
  - ➤ Saw tooth ease of subsequent harvesting of QC's
  - > 2-3 spikes 5-10 QC's
- Foundation or drawn comb
- Insert into parent colony
  - ➤ In centre of brood cluster
  - Check at 3 days intervals
  - Developing larvae in centre
  - ➤ Day old eggs on edge






- Set up Cell Raising Colony Simple Method
  - > Use a colony which is prone to swarming, often good at raising queens.
  - ➤ Utilise emergency response Remove queen.
  - Ensure ample pollen in adjacent frame and good income of nectar.
- Set up Cell Raising Colony Assembled from various colonies
  - ➤ Lots of young bees (shake in 6-8 frames into a 5/6 frame nuc)
  - Frames of capped or well developed larvae
  - Nectar or uncapped/unripe honey & frame of pollen
  - ➤ Gap left in middle for Breeding-Comb




- Trim Frame
  - ➤ Remove Eggs
  - ➤ Expose larvae 24-48hr old






- Trim Frame
  - ➤ Remove Eggs
  - Expose larvae 24-48hr old
- Insert into gap in starter colony






- Trim Frame
  - ➤ Remove Eggs
  - Expose larvae 24-48hr old
- Insert into gap in starter colony
- Inspect for Queen cells
  - ➤ 9 days on QC's capped
  - ➤ Larvae pupated
  - Easy to access / cut out






## Demaree Manipulation

Swarm Prevention – Brood Relocation







## Queen Mating

- Depending on the number of Queens needed, cell raising colony or Demaree top brood box could be broken down into 2 or 3 frame nucs.
- Mini mating nucs.
  - **≻**Apidea





## Queen Mating

- Depending on the number of Queens needed, cell raising colony or Demaree top brood box could be broken down into 2 or 3 frame nucs.
- Mini mating nucs.
  - ➤ Apidea Smallest
  - ➤ Kieler Small





## Queen Mating

- Depending on the number of Queens needed, cell raising colony or Demaree top brood box could be broken down into 2 or 3 frame nucs.
- Mini mating nucs.
  - ➤ Apidea Smallest
  - ➤ Kieler Small
  - ➤ Queen Castle
    - Full size frames





